



FEP Medical Policy Manual

FEP 2.01.58 Transanal Radiofrequency Treatment of Fecal Incontinence

Effective Policy Date: April 1, 2020

Related Policies:

None

Original Policy Date: June 2012

Transanal Radiofrequency Treatment of Fecal Incontinence

Description

Radiofrequency energy has been investigated as a minimally invasive treatment of fecal incontinence, in a procedure referred to as the Secca procedure. In this outpatient procedure using conscious sedation, radiofrequency energy is delivered to the sphincteric complex of the anal canal to create discrete thermal lesions. Over several months, these lesions heal and the tissue contracts, changing the tone of the tissue and improving continence.

OBJECTIVE

The objective of this evidence review is to determine whether transanal radiofrequency treatment for fecal incontinence improves the net health outcome compared with alternatives.

POLICY STATEMENT

Transanal radiofrequency therapy is considered **investigational** as a treatment of fecal incontinence.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

POLICY GUIDELINES

The Secca procedure may be performed on an outpatient basis using conscious sedation and a local anesthetic.

BENEFIT APPLICATION

Experimental or investigational procedures, treatments, drugs, or devices are not covered (See General Exclusion Section of brochure).

FDA REGULATORY STATUS

In 2002, the Secca™ System (Mederi Therapeutics) was cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process for "general use in the electrosurgical coagulation of tissue and is intended for use specifically in the treatment of fecal incontinence in those patients with incontinence to solid or liquid stool at least once per week and who have failed more conservative therapy."¹ Food and Drug Administration product code: GEI.

RATIONALE

Summary of Evidence

For individuals who have fecal incontinence who receive transanal radiofrequency (RF) treatment, the evidence includes 8 nonrandomized studies. Relevant outcomes are symptoms, change in disease status, quality of life, and treatment-related morbidity. Studies include a small number of patients, and estimates of treatment differences are very imprecise. Study follow-up periods vary and need to be considerably longer and involve larger numbers of patients to evaluate long-term outcomes properly. Three-year follow-up of a small cohort showed decrement in response over time. Multicenter randomized controlled trials with sufficient power are required to evaluate the continuing use of this procedure as an alternative to other surgical interventions, physical therapies, or as an adjunctive treatment option for fecal incontinence. The evidence is insufficient to determine the effects of the technology on health outcomes.

SUPPLEMENTAL INFORMATION

Practice Guidelines and Position Statements

National Institute for Health and Care Excellence

The National Institute for Health and Care Excellence (NICE) issued guidance on radiofrequency treatment for fecal incontinence in 2011.¹² NICE concluded that "evidence on endoscopic radiofrequency therapy of the anal sphincter for [fecal] incontinence raises no major safety concerns. There is evidence of efficacy in the short term but in a limited number of patients."

In 2016, NICE published a Medtech innovation briefing on the Secca system for fecal incontinence.¹³ These briefings aim to aid in the decision-making process by describing the technology, its role in the treatment pathway, the relevant published evidence, and cost information. These briefings do not contain recommendations. The briefing noted that "Secca therapy is a minimally invasive treatment option available for people with incontinence of solid or liquid stool at least once a week, in whom conservative management options have not controlled symptoms."

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

American Society of Colon and Rectal Surgeons

The American Society of Colon and Rectal Surgeons, in its 2015 clinical practice guidelines, noted: "Application of temperature-controlled radiofrequency energy to the sphincter complex may be used to treat fecal incontinence. Grade of Recommendation: Weak recommendation based on moderate-quality evidence, 2B."¹⁴ The guidelines also stated: "Because of the limitations in the available data, alternative treatments should be pursued before considering radiofrequency energy delivery."

American College of Gastroenterology

The American College of Gastroenterology published guidelines on the management of benign anorectal disorders in 2014.¹⁵ The guidelines indicated that there is insufficient evidence to recommend radiofrequency ablation to the anal sphincter as treatment for fecal incontinence. The College also asserted that the biologic rationale for this type of treatment is unproven.

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

REFERENCES

1. Food and Drug Administration (FDA). Attachment 14: 510(k) Summary. Curon Medical, Inc.'s Secca™ System. 2002; http://www.accessdata.fda.gov/cdrh_docs/pdf/k014216.pdf. Accessed September, 2019.
2. Forte ML, Andrade KE, Butler M, et al. Treatments for Fecal Incontinence (Comparative Effectiveness Review No. 165). Rockville, MD: Agency for Healthcare Research and Quality; 2016.
3. Abbas MA, Tam MS, Chun LJ. Radiofrequency treatment for fecal incontinence: is it effective long-term? *Dis Colon Rectum*. May 2012;55(5):605-610. PMID 22513440.
4. Rothbarth J, Bemelman WA, Meijerink WJ, et al. What is the impact of fecal incontinence on quality of life? *Dis Colon Rectum*. Jan 2001;44(1):67-71. PMID 11805565.
5. Efron JE, Corman ML, Fleshman J, et al. Safety and effectiveness of temperature-controlled radio-frequency energy delivery to the anal canal (Secca procedure) for the treatment of fecal incontinence. *Dis Colon Rectum*. Dec 2003;46(12):1606-1616; discussion 1616-1608. PMID 14668584.
6. Felt-Bersma RJ, Szojda MM, Mulder CJ. Temperature-controlled radiofrequency energy (SECCA) to the anal canal for the treatment of faecal incontinence offers moderate improvement. *Eur J Gastroenterol Hepatol*. Jul 2007;19(7):575-580. PMID 17556904.
7. Lam TJ, Visscher AP, Meurs-Szojda MM, et al. Clinical response and sustainability of treatment with temperature-controlled radiofrequency energy (Secca) in patients with faecal incontinence: 3 years follow-up. *Int J Colorectal Dis*. Jun 2014;29(6):755-761. PMID 24805249.
8. Ruiz D, Pinto RA, Hull TL, et al. Does the radiofrequency procedure for fecal incontinence improve quality of life and incontinence at 1-year follow-up? *Dis Colon Rectum*. Jul 2010;53(7):1041-1046. PMID 20551757.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.

9. Lefebure B, Tuech JJ, Bridoux V, et al. Temperature-controlled radio frequency energy delivery (Secca procedure) for the treatment of fecal incontinence: results of a prospective study. *Int J Colorectal Dis.* Oct 2008;23(10):993-997. PMID 18594840.
10. Takahashi-Monroy T, Morales M, Garcia-Osogobio S, et al. SECCA procedure for the treatment of fecal incontinence: results of five-year follow-up. *Dis Colon Rectum.* Mar 2008;51(3):355-359. PMID 18204954.
11. Kim DW, Yoon HM, Park JS, et al. Radiofrequency energy delivery to the anal canal: is it a promising new approach to the treatment of fecal incontinence? *Am J Surg.* Jan 2009;197(1):14-18. PMID 18614149.
12. National Institute for Health and Care Excellence (NICE). Endoscopic radiofrequency therapy of the anal sphincter for faecal incontinence [IPG393]. 2011; <https://www.nice.org.uk/guidance/ipg393>. Accessed September, 2019.
13. National Institute for Health and Care Excellence (NICE). Secca System for faecal incontinence [MIB66]. 2016; <https://www.nice.org.uk/advice/mib66/chapter/summary>. Accessed September, 2019.
14. Paquette IM, Varma MG, Kaiser AM, et al. The American Society of Colon and Rectal Surgeons' clinical practice guideline for the treatment of fecal incontinence. *Dis Colon Rectum.* Jul 2015;58(7):623-636. PMID 26200676.
15. Wald A, Bharucha AE, Cosman BC, et al. ACG clinical guideline: management of benign anorectal disorders. *Am J Gastroenterol.* Aug 2014;109(8):1141-1157; (Quiz) 1058. PMID 25022811.

POLICY HISTORY - THIS POLICY WAS APPROVED BY THE FEP® PHARMACY AND MEDICAL POLICY COMMITTEE ACCORDING TO THE HISTORY BELOW:

Date	Action	Description
June 2012	New Policy	
December 2013	Replace policy	Policy updated with literature review, no references added, no change in policy statement
December 2014	Replace policy	Policy updated with literature review; reference 6 added; no change in policy statement.
March 2016	Replace policy	Policy updated with literature review through November 10, 2015; reference 12 added. Policy statement unchanged.
March 2017	Replace policy	Policy updated with literature review through November 2, 2016; references 2, 13, and 15 added. Policy statement changed from not medically necessary to investigational.
March 2018	Replace policy	Policy updated with literature review through September 11, 2017; no references added. Policy statement unchanged.
March 2019	Replace policy	Policy updated with literature review through September 4, 2018; no references added. Policy statement unchanged.
March 2020	Replace policy	Policy updated with literature review through September 23, 2019; no references added. Policy statement unchanged.

The policies contained in the FEP Medical Policy Manual are developed to assist in administering contractual benefits and do not constitute medical advice. They are not intended to replace or substitute for the independent medical judgment of a practitioner or other health care professional in the treatment of an individual member. The Blue Cross and Blue Shield Association does not intend by the FEP Medical Policy Manual, or by any particular medical policy, to recommend, advocate, encourage or discourage any particular medical technologies. Medical decisions relative to medical technologies are to be made strictly by members/patients in consultation with their health care providers. The conclusion that a particular service or supply is medically necessary does not constitute a representation or warranty that the Blue Cross and Blue Shield Service Benefit Plan covers (or pays for) this service or supply for a particular member.